

The Paskalev-Tshobanov Distance Formula and Some of its Applications

SAVA GROZDEV^a, HIROSHI OKUMURA^b AND DEKO DEKOV^c ²

^a VUZF University of Finance, Business and Entrepreneurship,

Gusla Street 1, 1618 Sofia, Bulgaria

e-mail: sava.grozdev@gmail.com

^b Maebashi Gunma, 371-0123, Japan

e-mail: hokmr@yandex.com

^cZahari Knjazheski 81, 6000 Stara Zagora, Bulgaria

e-mail: ddekov@ddekov.eu

web: <http://www.ddekov.eu/>

Abstract. By using the Paskalev-Tchobanov distance formula, we give new proofs of a few well-known formulas from Euclidean geometry.

Keywords. Euclidean geometry, triangle geometry, Paskalev-Tchobanov distance formula.

Mathematics Subject Classification (2010). 51-04, 68T01, 68T99.

The Paskalev-Tchobanov distance formula, published in 1985 (See §15, [2] and formula (9) in [1]), is a powerful tool for finding distances in triangle geometry.

By using the Paskalev-Tchobanov distance formula, we will give new proofs of a few well-known formulas from triangle geometry. There is a large number of formulas in triangle geometry which could be easily proved by the approach of this paper.

We encourage the reader to extend the list of Theorems 2-7 given in this paper. .

Given triangle ABC with side-lengths $a = BC, b = CA, c = AB$. We denote by Δ the area of triangle ABC . Denote by

- $s = \frac{a + b + c}{2}$ the semiperimeter of triangle ABC .
- Δ the area of triangle ABC .
- r the radius of the Incircle.
- R the radius of the Circumcircle.

¹This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution, and reproduction in any medium, provided the original author(s) and the source are credited.

²Corresponding author

- $m_a = AM_a$, $m_b = BM_b$, $m_c = CM_c$, the medians.
- $l_a = AL_a$, $l_b = BL_b$, $l_c = CL_c$, the internal bisectors.
- $h_a = AH_a$, $h_b = BH_b$, $h_c = CH_c$, the altitudes.

Denote by

- I the Incenter.
- O the Circumcenter.
- G_a projections of the Incenter on side BC .

The barycentric coordinates of vertices of triangle ABC are $A = (1, 0, 0)$, $B = (0, 1, 0)$, $C = (0, 0, 1)$. The barycentric coordinates of the following points are well-known (See e.g. [1], [3]):

$$\begin{aligned} I &= (a, b, c), \quad O = (a^2(b^2 + c^2 - a^2), b^2(c^2 + a^2 - b^2), c^2(a^2 + b^2 - c^2)), \\ M_a &= (0, 1, 1), \quad M_b = (1, 0, 1), \quad M_c = (1, 1, 0), \\ L_a &= (0, b, c), \quad L_b = (a, 0, c), \quad L_c = (a, b, 0), \\ H_a &= (0, (a^2 + b^2 - c^2)(b^2 + c^2 - a^2), (b^2 + c^2 - a^2)(c^2 + a^2 - b^2)), \\ G_a &= (0, (b - c + a)(b + c - a), (c - a + b)(c + a - b)). \end{aligned}$$

The Paskalev-Tcobanov distance theorem states:

Theorem 1. *Given two points $P = (u_1, v_1, w_1)$ and $Q = (u_2, v_2, w_2)$ in normalized barycentric coordinates. Denote $x = u_1 - u_2$, $y = v_1 - v_2$ and $z = w_1 - w_2$. Then the square of the distance between P and Q is*

$$PQ^2 = -a^2yz - b^2zx - c^2xy.$$

Theorem 2. $\Delta = \sqrt{s(s-a)(s-b)(s-c)}$. (Heron's Formula)

Proof. By using Theorem 1, we find the length of the altitude

$$h_a = AH_a = \frac{2\sqrt{s(s-a)(s-b)(s-c)}}{a}.$$

Hence $\Delta = \frac{ah_a}{2} = \sqrt{s(s-a)(s-b)(s-c)}$ □

Theorem 3. $R = \frac{abc}{4\Delta}$.

Proof. By using Theorem 1, we obtain $R = AO = \frac{abc}{4\sqrt{s(s-a)(s-b)(s-c)}}$, and

by using Herons's formula we obtain $R = \frac{abc}{4\Delta}$, □

Theorem 4. $r = \frac{\Delta}{s}$

Proof. By using Theorem 1, we obtain $r = IG_a = \frac{\sqrt{s(s-a)(s-b)(s-c)}}{s}$, and

by using the Herons's formula we obtain $r = \frac{\Delta}{s}$. □

Theorem 5.

$$m_a = \frac{\sqrt{2b^2 + 2c^2 - a^2}}{2}, \quad m_b = \frac{\sqrt{2c^2 + 2a^2 - b^2}}{2}, \quad m_c = \frac{\sqrt{2a^2 + 2b^2 - c^2}}{2}.$$

Proof. By using Theorem 1, we obtain $m_a = AM_a = \frac{\sqrt{2b^2 + 2c^2 - a^2}}{2}$. Similarly for m_b and m_c . \square

Theorem 6. $l_a = \frac{\sqrt{bcs(s-a)}}{b+c}$, $l_b = \frac{\sqrt{cas(s-b)}}{c+a}$, $l_c = \frac{\sqrt{abs(s-c)}}{a+b}$.

Proof. By using Theorem 1, we obtain $l_a = AL_a = \frac{\sqrt{bcs(s-a)}}{b+c}$. Similarly for l_b and l_c . \square

Theorem 7. $OI^2 = R(R - 2r)$. (*Euler formula*)

Proof. By using Theorems 3 and 4 we obtain $R(R - 2r) = \frac{abcE}{16\Delta^2}$, where

$$E = a^3 + b^3 + c^3 + 3abc - ba^2 - b^2a - ca^2 - c^2a - c^2b - b^2c.$$

By using Theorem 1, we obtain $OI^2 = \frac{abcE}{16\Delta^2}$. Hence $OI^2 = R(R - r)$. \square

REFERENCES

- [1] S. Grozdev and D. Dekov, Barycentric Coordinates: Formula Sheet, International Journal of Computer Discovered Mathematics, vol.1, 2016, no 2, 75-82. <http://www.journal-1.eu/2016-2/Grozdev-Dekov-Barycentric-Coordinates-pp.75-82.pdf>.
- [2] G. Paskalev and I. Tchobanov, *Remarkable Points in the Triangle* (in Bulgarian), Sofia, Narodna Prosveta, 1985.
- [3] E. W. Weisstein, *MathWorld - A Wolfram Web Resource*, <http://mathworld.wolfram.com/>.